



# Radioactive Concentrations in Chemical Fertilizers

Gwang-Ho Kim<sup>1</sup>, Jae-Hwan Cho<sup>2</sup>

<sup>1</sup>Department of Medical Device Management and Research, Samsung Advanced Institute for Health Sciences & Technology, Sungkyunkwan University, Seoul, Korea; <sup>2</sup>Department of Radiologic Science, Ansan University, Ansan, Korea

## ABSTRACT

**Background:** The aim of the present study was to determine radioactive concentrations in fertilizers known to contain essential nutrients. Results of this study could be used as basic data to monitor the impact of chemical fertilizers on the environment and public health. Nitrogen fertilizers, calcium fertilizers, sulfur fertilizers, phosphate acid fertilizers, and potassium chloride fertilizers were used in this study.

**Materials and Methods:** Five chemical fertilizers were pulverized, placed in polyethylene containers, and weighed. The time to measure each specimen was set to be 3,600 seconds for a scintillator-based gamma-ray spectroscopy system. Concentration of gamma radionuclide was analyzed based on obtained spectra. At the end of the measurement, the spectrum file was stored and used to calculate radioactive concentrations using a gamma-ray spectrometer software.

**Results and Discussion:** In the nitrogen fertilizer,  $3.49 \pm 5.71$  Bq/kg of <sup>137</sup>Cs,  $34.43 \pm 7.61$  Bq/kg of <sup>134</sup>Cs, and  $569.16 \pm 91.15$  of <sup>40</sup>K were detected whereas <sup>131</sup>I was not detected. In the calcium fertilizer,  $5.74 \pm 4.40$  Bq/kg of <sup>137</sup>Cs (the highest concentration among all fertilizers),  $22.37 \pm 5.39$  Bq/kg of <sup>134</sup>Cs, and  $433.67 \pm 64.24$  Bq/kg of <sup>40</sup>K were detected whereas <sup>131</sup>I was not detected. In the sulfur fertilizer,  $347.31 \pm 55.73$  Bq/kg of <sup>40</sup>K,  $19.42 \pm 4.53$  Bq/kg of <sup>134</sup>Cs,  $2.21 \pm 3.49$  of <sup>137</sup>Cs, and  $0.04 \pm 0.22$  Bq/Kg of <sup>131</sup>I were detected. In the phosphoric acid fertilizer,  $70,007.34 \pm 844.18$  Bq/kg of <sup>40</sup>K (the highest concentration among all fertilizers) and  $46.07 \pm 70.40$  Bq/kg of <sup>134</sup>Cs were detected whereas neither <sup>137</sup>Cs nor <sup>131</sup>I was detected. In the potassium chloride fertilizer,  $12,827.92 \pm 1542.19$  Bq/kg of <sup>40</sup>K was and  $94.76 \pm 128.79$  Bq/kg of <sup>134</sup>Cs were detected whereas neither <sup>137</sup>Cs nor <sup>131</sup>I was detected. The present study examined inorganic fertilizers produced by a single manufacturer. There might be different results according to the country and area from which fertilizers are imported. Further studies about inorganic fertilizers in more detail are needed to create measures to reduce <sup>40</sup>K.

**Conclusion:** Measures are needed to reduce radiation exposure to <sup>40</sup>K contained in fertilizers including phosphoric acid and potassium chloride fertilizers.

**Keywords:** Radioactive concentration, Fertilizers, Scintillator

## Original Research

**Received** July 7, 2021

**Revision** October 15, 2021

**Accepted** August 1, 2022

**Corresponding author:** Jae-Hwan Cho

Department of Radiologic Science, Ansan University, 155 Ansan University-ro, Sangnok-gu, Ansan 15328, Korea  
E-mail: 8452404@hanmail.net

<https://orcid.org/0000-0001-7112-3866>

This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by-nc/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright © 2022 The Korean Association for Radiation Protection

## Introduction

Ionizing radiation was discovered by German physicist Wilhelm Conrad Rontgen in 1895. Since then, radiation has been used in various industries such as manufacturing, agriculture, health, and high technologies [1]. Accordingly, more and more industry sectors are using radiation after understanding its characteristics and energy properties. Radiation technologies are being more widely used in the area of research and development for daily necessities, including the health industry, advanced parts manufacturing, biotechnology, and new variety development [2]. With the use of radiation,

the quality of life has been improved along with development of industries and economies [3]. On average, our radiation exposure due to all natural sources amounts to about 2.4 mSv a year [4]. It is considered that natural radiation has a minimal effect on the human body. However, the International Commission on Radiological Protection has warned that risks of radiation exposure are likely to increase proportionately with higher exposure to radiation even in small doses [5]. It has also been reported that plants growing in a soil rich in natural radionuclides coming from uranium or thorium can play a part in internal exposure to radioactive elements that we take into our bodies [6]. Phosphate rock, a pre-concentrated phosphate ore, is the primary raw material for the production of mineral phosphate fertilizer. It contains 0.02% of uranium. In its by-product, phospho-gypsum, radioactive elements including radium exist. Caesium-137 is one of the radionuclides from various human activities, such as the spread of nuclear materials, for some reasons. It can cause food contamination and internal exposure to radiation [7]. After the Fukushima Daiichi accident in 2011, massive efforts have been put together to regulate and decrease  $^{137}\text{Cs}$  in food items in Japan [8–10]. There are pertinent risks of carcinogenesis related to internal radiation doses due to food contaminated by radioactive materials including  $^{134}\text{Cs}$ ,  $^{137}\text{Cs}$ , and  $^{131}\text{I}$  according to Korea National Health and Nutrition Examination Survey. Nuclear-fuel reprocessing plants, nuclear wastes, and geological repositories are main sources of radioactive iodine in the environment [11, 12]. Conventionally, radiation exposures to uranium, radium, thorium, and potassium are well accounted for due to their roles in radioactivity from the environment. However, after recent accidents such as the Fukushima Daiichi accident in 2011, it is now of dire importance to measure radiation levels in South-east Asia generally and Korea specifically due to its very close proximity to Fukushima, Japan [13, 14]. Iodine-131 is a very critical radioisotope which is closely associated with nuclear fission reactions. It causes environmental hazards in case of nuclear bomb attacks/mishaps and nuclear power plant accidents [15]. In case of a disaster or incident involving the spread of nuclear waste or a nuclear attack, the resultant radioactivity can spread up to 300 miles. The most extravagant health hazard from exposure to  $^{131}\text{I}$  is an enhanced risk of cancer induced due to radiation. It can also result in deformities, non-cancerous growths, and thyroiditis [16]. Accidents such as nuclear-reactor incidents at Chernobyl in 1986, the incident of Three Mile Island in 1979, and the most re-

cent Fukushima nuclear reactor accident in 2011 provide clear evidence that it is urgent to understand health hazards posed by the presence of iodine in the atmosphere [17]. After the Fukushima incident, Korean authorities are taking threats of food-based radiation very seriously. The Ministry of Food and Drug Safety in Korea has conducted surveys to measure radioactivity levels of  $^{134}\text{Cs}$ ,  $^{137}\text{Cs}$ , and  $^{131}\text{I}$  through the Korea National Health and Nutrition Examination Survey [12]. Geographically, Korea is the closest country to Japan. Therefore, there is a dire need to confirm the absence or presence of  $^{131}\text{I}$  in the environment and soil. Although the quantity of radiation by each radioactive element individually can be below the standard hazardous level or the maximum tolerance of 370 Bq/kg for any food, fertilizers with multiple radioactive elements used for farming might deliver radiation to agricultural products and consequently contaminate water reserves such as streams and groundwater after mixing with rain or irrigation water [6, 18–22]. Although there are a number of studies available about radioactivity due to  $^{238}\text{U}$ ,  $^{232}\text{Th}$ , and  $^{40}\text{K}$  present in chemical fertilizers from countries such as Serbia, Bangladesh, India, and Iraq [23–26], studies that measure the radioactivity of agricultural fertilizers in Korea and the rest of the world are lacking. Studies generally involve  $^{137}\text{Cs}$  and  $^{130}\text{I}$  specifically. A variety of reports have analyzed radioactivity levels in water, air, and soil [27–29]. Hence, this study will serve as a foundation to better monitor the impact of radioactive levels in chemical fertilizers on public health and environment and to ensure that radiation levels in chemical fertilizers that contain essential nutrients for plant growth and development are safe for human health and the environment.

## Materials and Methods

### 1. Experimental Materials

A total of five fertilizers, nitrogen fertilizers, calcium fertilizers, sulfur fertilizers, phosphate acid fertilizers, and potassium chloride fertilizers, were used as experimental fertilizers. They are the most widely used fertilizers in Korea. Distilled water was used as a control (Fig. 1). Nitrogen fertilizers contained 46 % of nitrogen. Calcium fertilizers had 60% of calcium. Sulfur fertilizers had 24% of sulfur. Phosphate acid fertilizers had 50% of soluble phosphate acid, 32% of soluble chloride, 0.05% of soluble copper, and 0.05% of soluble zinc. Potassium chloride fertilizers had 60% of potassium. Fertilizers are named with the highest content of the main ingredient.

## 2. Methods

### 1) Sample preparation

Five specimens (nitrogen fertilizer, calcium fertilizer, sulfur fertilizer, phosphate acid fertilizer, and potassium chloride fertilizer) and distilled water as control were used (Table 1). Each of these five specimens was pulverized and placed in a 500 mL of Marinelli vessel (Fig. 2). The density of the distilled



**Fig. 1.** Distilled water (A), nitrogen (B), calcium (C), sulfur (D), phosphate acid (E), and potassium chloride (F) samples used in this study.

**Table 1.** Specimen Information

Classification	Fertilizer type	Weight (g)	Volume (mL)
a	Water	500	500
b	Nitrogen	548	500
c	Calcium	414	500
d	Sulfur	893	500
e	Phosphate acid	845	500
f	Potassium chloride	873	500

**Table 2.** Composition of Distilled Water

Category	General constants (mg/L)
Fluoride ion	0.34
Chlorine ion	10.7
Nitric nitrogen	1.30
Sulfuric acid ion	11.1
Aluminum	0.011
Copper	0.035
Iron	0.219

water ranged from 0.998 g/cm<sup>3</sup> at 20 °C to 0.996 g/cm<sup>3</sup> at 30 °C. Compositions of distilled water are shown in Table 2.

Each container was weighed on a scale. This time, the container weight was not considered. The net weight was marked on each container and numbered. The distilled water and each specimen were placed on a thallium-activated sodium iodide, NaI(Tl), scintillator-based gamma-ray spectroscopy



**Fig. 2.** Chemical fertilizers in scintillator. An empty container as a background (A), nitrogen (B), calcium (C), sulfur (D), phosphate acid (E), and potassium chloride (F) fertilizers are shown.

**Table 3.** Characteristics of the NaI Detector

Variable	Value
Detector unit	NaI(Tl) Ø 40 mm × 40 mm
Number of channels	1,024
Energy range	0–3 MeV
Background	<40 counts per minute (cpm)
Measurable activity concentration range of <sup>137</sup> Cs	5–1,000,000 Bq/kg
Measurable activity concentration range of <sup>134</sup> Cs	5–1,000,000 Bq/kg
Measurable activity concentration range of <sup>40</sup> K	50–20,000 Bq/kg
Intrinsic measurement error at P=0.95	±20%
Dimensions	255 mm × 360 mm × 300 mm

system (gamma-ray spectrometer RUG 91-2; LINEV Systems, Minsk, Belarus) and measured. External radiation was blocked with a shield consisting of 48% lead, 45% tungsten, 5% iron, and 2% nickel with a thickness of 10 cm. Characteristics of the NaI(Tl) detector are provided in Table 3. The NaI(Tl) detector can respond to a gamma ray by producing a small flash of light or a scintillation. The scintillation occurs when scintillator electrons excited by energy of photon return to their ground state. The detector crystal is mounted on a photomultiplier tube which converts scintillation into an electrical pulse. This is taken from the anode of the photomultiplier. It is a negative pulse. The NaI(Tl) crystal is protected from the moisture in the air by encasing it in aluminum, which also serves as a convenient mounting for the entire crystal/photomultiplier unit.

## 2) Spectrometric analysis

RUG 91-2 gamma-ray spectrometer (radiometer) was used to analyze the amplitude of photon distribution of pulses generated from the scintillation detector upon detection of gamma-ray of the sample. Amplitude distribution of pulses was then analyzed and radionuclide activity was calculated. In order to enhance the registration of photon impulses, the sample was placed into a 0.5-L Marinelli vessel installed in a lead shield which could protect the sample from the influence of an external background radiation. Calibration and check took only a few minutes. Photo peaks of gamma-ray at 1461 keV, 622 keV, 796 keV, and 365 keV were used for measuring  $^{40}\text{K}$ ,  $^{137}\text{Cs}$ ,  $^{134}\text{Cs}$ , and  $^{131}\text{I}$ , respectively.

Distilled water as control and each fertilizer specimen were counted for 3,600 seconds. Gamma radionuclide concentration was then analyzed to obtain a spectrum file. At the end of each measurement, the spectrum file which had been stored was analyzed using a gamma-ray spectrometer software. Based on the analysis results, radiation level of each specimen was then determined.

## Results and Discussion

The nitrogen fertilizer had  $3.49 \pm 5.71$  Bq/kg of  $^{137}\text{Cs}$ , which was the second highest concentration,  $34.43 \pm 7.61$  Bq/kg of  $^{134}\text{Cs}$ , and  $569.16 \pm 91.15$  Bq/kg of  $^{40}\text{K}$ , whereas  $^{131}\text{I}$  was not detected. The calcium fertilizer had the highest concentration of  $^{137}\text{Cs}$  at  $5.74 \pm 4.40$  Bq/kg. It also had  $22.37 \pm 5.39$  Bq/kg of  $^{134}\text{Cs}$  and  $433.67 \pm 64.24$  Bq/kg of  $^{40}\text{K}$ . However,  $^{131}\text{I}$  was not detected. The sulfur fertilizer had  $347.31 \pm 55.73$  Bq/kg of  $^{40}\text{K}$ ,  $19.42 \pm 4.53$  Bq/kg of  $^{134}\text{Cs}$ ,  $2.21 \pm 3.49$  Bq/kg of  $^{137}\text{Cs}$ , and  $0.04 \pm 0.22$  Bq/kg for  $^{131}\text{I}$ . The phosphoric acid fertilizer had the highest concentration of  $^{40}\text{K}$  at  $70,007.34 \pm 844.18$  Bq/kg and  $46.07 \pm 70.40$  Bq/kg of  $^{134}\text{Cs}$ , whereas neither  $^{137}\text{Cs}$  nor  $^{131}\text{I}$  was detected. The potassium chloride fertilizer had  $12,827.92 \pm 1,542.19$  Bq/kg of  $^{40}\text{K}$  and  $94.76 \pm 128.79$  Bq/kg of  $^{134}\text{Cs}$ . Neither  $^{137}\text{Cs}$  nor  $^{131}\text{I}$  was detected (Table 4). Fig. 3 shows radioactive concentrations and spectra of the five fertilizers and distilled water measured for 3,600 seconds.

Currently, radioactive substances manufactured or processed using radiation technologies are being accumulated worldwide. They are mostly radioactive wastes produced from nuclear power plants, fallouts from nuclear tests, and radioisotopes used in the health industry [30]. Due to some nuclear disasters such as the Chernobyl nuclear disaster in 1986 and the Russia nuclear disaster in 1993, people are concerned about impacts of radiation on human health [31]. As many people are exposed to radionuclides from natural radiation sources in the soil due to such accidents, it is important to minimize radiation impacts. Zmazek et al. [32] have reported that areas rich in granite are more likely to have uranium and radon than non-granite areas. Granites include 2–12 parts per million (ppm) of uranium and 8–33 ppm thorium [33]. Every year, the Korea Institute of Nuclear Safety measures radiation in subsoil and topsoil for 14 areas and publishes analysis results due to concern about radiation impacts. Thus, the objective of this study was to measure ra-

**Table 4.** Activity Concentration of  $^{134}\text{Cs}$ ,  $^{137}\text{Cs}$ ,  $^{40}\text{K}$ , and  $^{131}\text{I}$  Radionuclides in Samples

Sample	Fertilizer	Activity concentration (Bq/kg)			
		$^{134}\text{Cs}$	$^{137}\text{Cs}$	$^{40}\text{K}$	$^{131}\text{I}$
b	Nitrogen	$34.43 \pm 7.61$	$3.49 \pm 5.71$	$569.16 \pm 91.15$	<LLD
c	Calcium	$22.37 \pm 5.39$	$5.74 \pm 4.40$	$433.67 \pm 64.24$	<LLD
d	Sulfur	$19.42 \pm 4.53$	$2.21 \pm 3.49$	$347.31 \pm 55.73$	$0.04 \pm 0.22$
e	Phosphate acid	$46.07 \pm 70.40$	<LLD	$70,007.34 \pm 844.18$	<LLD
f	Potassium chloride	$94.76 \pm 128.79$	<LLD	$12,827.92 \pm 1,542.19$	<LLD

Values are presented as mean  $\pm$  standard deviation.  
LLD, lower limit of detection.

diation levels of inorganic fertilizers widely used in agriculture. As a result,  $^{40}\text{K}$  showed the highest levels in fertilizers containing phosphate acid and potassium chloride. A phosphate acid fertilizer's base material is phosphate rock, a nat-

ural radiation source which contains  $^{235}\text{U}$ ,  $^{238}\text{U}$ , and  $^{232}\text{Th}$ . Phosphate rock exporters to Korea are Morocco, China, Israel, and Togo. It was found that rocks they exported exceeded the radiation standard (1,000 Bq/kg) [34] in terms of  $^{235}\text{U}$  and




Fig. 3. Radioactive concentration spectrum of each fertilizer measured for 3,600 seconds. Spectra of background (A), nitrogen (B), calcium (C), sulfur (D), phosphate acid (E), and potassium chloride (F) are shown. (Continued to the next page)




Fig. 3. Continued.

$^{238}\text{U}$ . As potassium chloride is added in order to improve hygroscopic capacity in the process of manufacturing a phosphate acid fertilizer, the level of  $^{40}\text{K}$  increases accordingly [35]. A potassium chloride fertilizer is manufactured with leucite, alunite, and sericite known to be rich in potassium. It has been reported that ores rich in potassium imported by Korea from countries such as Canada, Belarus, and Russia have high  $^{40}\text{K}$  levels, exceeding 1,000 Bq/kg. Almost all foods

have natural radioactivity due to radioactive isotopes. For example,  $^{40}\text{K}$  is present at 40–50 Bq/kg in cows' milk, 400–500 Bq/kg in milk powder, 600–800 Bq/kg in concentrated fruit juice, and over 1000 Bq/kg in instant coffee [22]. Natural radioactivity is primarily due to primordial radionuclides  $^{40}\text{K}$ ,  $^{232}\text{Th}$ ,  $^{238}\text{U}$ ,  $^{235}\text{U}$ , and so on present in different quantities depending on soils of different regions in the world [36]. According to the International Atomic Energy Agency, a per-

missible limit for food is 370 Bq/kg of  $^{40}\text{K}$  [35]. However, Rajacic et al. [25] have demonstrated that the concentration of  $^{40}\text{K}$  in fertilizer is 4,860 Bq/kg [25]. Radiation due to  $^{40}\text{K}$ , a radionuclide, is rapidly absorbed into the body, causing internal exposure to radioactive elements that we take into our bodies through food and water [37]. In the environment,  $^{40}\text{K}$  is redistributed through agriculture by application of fertilizers. Thus, its concentration should be measured to distinguish safe utilization levels of fertilizers rich in  $^{40}\text{K}$  [19, 25, 38]. From production to utilization in agriculture, fertilizers might cause exposure to gamma radiation (external exposure) [39–41]. Many biomedical hazards are associated with exposure to radiation for a few weeks or months to years, which might cause radiation-induced liver disease [42–45]. Exposure to a significant level of radiation released from fertilizers can potentially cause cancers. Thus, a deeper understanding of fertilizers-based radiation is required [46, 47]. In particular,  $^{40}\text{K}$  is a constant and uniform source of exposure to radiation for humans [48]. The present study examined inorganic fertilizers produced by a single manufacturer. Results might be different depending on the country and areas from which fertilizers are imported. Further studies about inorganic fertilizers in more details are needed to create measures to reduce  $^{40}\text{K}$ .

## Conclusion

As a result of this study,  $^{40}\text{K}$  had the highest levels in five fertilizers. Thus, there is a need to take measures to reduce radiation exposure from  $^{40}\text{K}$  contained in fertilizers including phosphoric acid and potassium chloride. This study can help us understand radioactive concentrations in five inorganic fertilizers used in Korea. In the future, it is necessary to set up measures to cut radiation impacts from radionuclides.

## Conflict of Interest

No potential conflict of interest relevant to this article was reported.

## Ethical Statement

This article does not contain any studies with human participants or animals performed by any of the authors.

## Author Contribution

Conceptualization: Kim GH. Project administration: Cho JH. Writing - original draft: Kim GH. Writing - review and editing: Cho JH. Approval of final manuscript: all authors.

## References

1. Glasser O. W. C. Roentgen and the discovery of the Roentgen rays. *AJR Am J Roentgenol*. 1995;165(5):1033–1040.
2. Byun MW, Yook HS. Internal and external situation of irradiation technology utilization in the food and public health industry. *Korean J Food Preserv*. 2003;10(1):106–123.
3. Heo J. Radiation biology. Seoul: Shinkwang Publishing; 1998.
4. Mettler FA, Sinclair WK, Anspaugh L, Edington C, Harley JH, Ricks RC, et al. The 1986 and 1988 UNSCEAR (United Nations Scientific Committee on the Effects of Atomic Radiation) reports: findings and implications. *Health Phys*. 1990;58(3):241–250.
5. International Commission on Radiological Protection. 1990 Recommendations of the International Commission on Radiological Protection (ICRP Publication 60). *Ann ICRP*. 1991;21(1–3): 1–201.
6. Absalom JP, Young SD, Crout NM, Sanchez A, Wright SM, Smolders E, et al. Predicting the transfer of radiocaesium from organic soils to plants using soil characteristics. *J Environ Radioact*. 2001;52(1):31–43.
7. Topping CE, Abella MK, Berkowitz ME, Molina MR, Nikolic-Hughes I, Hughes EW, et al. In situ measurement of cesium-137 contamination in fruits from the northern Marshall Islands. *Proc Natl Acad Sci U S A*. 2019;116(31):15414–15419.
8. Merz S, Shozugawa K, Steinhauser G. Analysis of Japanese radionuclide monitoring data of food before and after the Fukushima nuclear accident. *Environ Sci Technol*. 2015;49(5):2875–2885.
9. Hamada N, Ogino H, Fujimichi Y. Safety regulations of food and water implemented in the first year following the Fukushima nuclear accident. *J Radiat Res*. 2012;53(5):641–671.
10. Hamada N, Ogino H. Food safety regulations: what we learned from the Fukushima nuclear accident. *J Environ Radioact*. 2012; 111:83–99.
11. Francis AJ, Nancharaiah YV. In situ and ex situ bioremediation of radionuclide-contaminated soils at nuclear and norm sites. In: Van Velzen L, editor. Environmental remediation and restoration of contaminated nuclear and norm sites. Sawston, UK: Woodhead Publishing; 2015. p. 185–236.
12. Moon EK, Ha WH, Seo S, Jin YW, Jeong KH, Yoon HJ, et al. Estimates of radiation doses and cancer risk from food intake in Korea. *J Korean Med Sci*. 2016;31(1):9–12.
13. Schnug E, Lottermoser BG. Fertilizer-derived uranium and its

- threat to human health. *Environ Sci Technol.* 2013;47(6):2433–2434.
14. Khater AE, AL-Sewaidan HA. Radiation exposure due to agricultural uses of phosphate fertilizers. *Radiat Meas.* 2008;43(8):1402–1407.
  15. Fuge R. Iodine in the environment its distribution and relationship to human health [Internet]. Vienna, Austria: International Atomic Energy Agency; 1987 [cited 2022 Jan 4]. Available from: <https://inis.iaea.org/search/searchsinglerecord.aspx?recordsFor=SingleRecord&RN=19105067>.
  16. Rivkees SA, Sklar C, Freemark M. Clinical review 99: the management of Graves' disease in children, with special emphasis on radioiodine treatment. *J Clin Endocrinol Metab.* 1998;83(11):3767–3776.
  17. Braverman ER, Blum K, Loeffke B, Baker R, Kreuk F, Yang SP, et al. Managing terrorism or accidental nuclear errors, preparing for iodine-131 emergencies: a comprehensive review. *Int J Environ Res Public Health.* 2014;11(4):4158–4200.
  18. Kadi MW, Al-Eryani DA. Natural radioactivity and radon exhalation in phosphate fertilizers. *Arab J Sci Eng.* 2012;37(1):225–231.
  19. International Atomic Energy Agency. Measurement of radionuclides in food and the environment: a guidebook. Vienna, Austria: International Atomic Energy Agency; 1989.
  20. Marovic G, Sencar J. <sup>226</sup>Ra and possible water contamination due to phosphate fertilizer production. *J Radioanal Nucl Chem.* 1995;200(1):9–18.
  21. Righi S, Lucialli P, Bruzzi L. Health and environmental impacts of a fertilizer plant: part I: assessment of radioactive pollution. *J Environ Radioact.* 2005;82(2):167–182.
  22. Stansfield CM. Legislation: contaminants and adulterants. In: Caballero B, editor. *Encyclopedia of food sciences and nutrition.* New York, NY: Academic Press; 2003. p. 3507–3513.
  23. Alam MN, Chowdhury MI, Kamal M, Ghose S, Banu H, Chakraborty D. Radioactivity in chemical fertilizers used in Bangladesh. *Appl Radiat Isot.* 1997;48(8):1165–1168.
  24. Ghosh D, Deb A, Bera S, Sengupta R, Patra KK. Measurement of natural radioactivity in chemical fertilizer and agricultural soil: evidence of high alpha activity. *Environ Geochem Health.* 2008;30(1):79–86.
  25. Rajacic MM, Sarap NB, Jankovic MM, Nikolic DJ, Todorovic DJ, Pantelic GK. Radioactivity in chemical fertilizers. Island of Krk, Croatia: Proceedings of 9<sup>th</sup> Symposium of the Croatian Radiation Protection Association; 2013 April 10–12.
  26. Azeez HH, Mansour HH, Ahmad ST. Effect of using chemical fertilizers on natural radioactivity levels in agricultural soil in the Iraqi Kurdistan region. *Pol J Environ Stud.* 2020;29(2):1059–1068.
  27. Peralta AA, Link MS, Schwartz J, Luttmann-Gibson H, Dockery DW, Blomberg A, et al. Exposure to air pollution and particle radioactivity with the risk of ventricular arrhythmias. *Circulation.* 2020;142(9):858–867.
  28. Bonotto DM, Bueno TO, Tessari BW, Silva A. The natural radioactivity in water by gross alpha and beta measurements. *Radiat Meas.* 2009;44(1):92–101.
  29. Faheem M, Mujahid SA, Matiullah. Assessment of radiological hazards due to the natural radioactivity in soil and building material samples collected from six districts of the Punjab province-Pakistan. *Radiat Meas.* 2008;43(8):1443–1447.
  30. Ulsh B, Hinton TG, Congdon JD, Dugan LC, Whicker FW, Bedford JS. Environmental biodosimetry: a biologically relevant tool for ecological risk assessment and biomonitoring. *J Environ Radioact.* 2003;66(1–2):121–139.
  31. Vandecasteele CM. Environmental monitoring and radioecology: a necessary synergy. *J Environ Radioact.* 2004;72(1–2):17–23.
  32. Zmazek B, Vaupotic J. Coping with radon problem in a private house. *Build Environ.* 2007;42(10):3685–3690.
  33. Lee HM, Moon KH, Kim JS, Ahn JK, Kim HC. Distribution of some environmental radionuclides in rocks and soils of Guemjeong-gu area in Busan, Korea. *J Petrol Soc Korea.* 2008;17(3):179–190.
  34. Ministry Law Korea. Hazardous limits of radiation [Internet]. Sejong: Korean Law Information Center; 2016 [cited 2022 Jan 4]. Available from: [https://www.law.go.kr/행정규칙/방사선방호등에관한기준/\(2016-16,20160831\)](https://www.law.go.kr/행정규칙/방사선방호등에관한기준/(2016-16,20160831)).
  35. Jang JG. Investigation and analysis of actual state of safety management for radiation in the natural environment. Seoul, Korea: Korean Industrial Hygiene Association; 2014.
  36. Missimer TM, Teaf C, Maliva RG, Danley-Thomson A, Covert D, Hegy M. Natural radiation in the rocks, soils, and groundwater of Southern Florida with a discussion on potential health impacts. *Int J Environ Res Public Health.* 2019;16(10):1793.
  37. Choi MS, Lin XJ, Lee SA, Kim W, Kang HD, Doh SH, et al. Daily intakes of naturally occurring radioisotopes in typical Korean foods. *J Environ Radioact.* 2008;99(8):1319–1323.
  38. International Atomic Energy Agency (IAEA). International basic safety standards for protection against ionizing radiation and for the safety of radiation sources [Internet]. Vienna: IAEA; 1996 [cited 2022 Jan 4]. Available from: <http://www-ns.iaea.org/standards/>
  39. Hassan NM, Mansour NA, Fayed-Hassan M, Fares S. Assessment of radiation hazards due to exposure to radionuclides in marble and ceramic commonly used as decorative building materials in Egypt. *Indoor Built Environ.* 2017;26(3):317–326.
  40. Hassan NM. Radon emanation coefficient and its exhalation rate of wasted petroleum samples associated with petroleum industry in Egypt. *J Radioanal Nucl Chem.* 2014;299(1):111–117.
  41. Iwaoka K, Tabe H, Yonehara H. Natural radioactivity of bedrock bath instruments and hot spring instruments in Japan. *J Radioanal Nucl Chem.* 2013;295(2):817–821.
  42. Bracht LK, Wen P, Meyerhardt JA, Kulke MH, Hornick JL, Redston M, et al. DNA repair enzyme expression and differential response



- to temozolomide in a patient with both glioblastoma and metastatic pancreatic neuroendocrine tumor. *J Clin Oncol*. 2008;26(29):4843–4844.
43. Lawrence TS, Robertson JM, Anscher MS, Jirtle RL, Ensminger WD, Fajardo LF. Hepatic toxicity resulting from cancer treatment. *Int J Radiat Oncol Biol Phys*. 1995;31(5):1237–1248.
44. Kim J, Jung Y. Radiation-induced liver disease: current understanding and future perspectives. *Exp Mol Med*. 2017;49(7):e359.
45. Benson R, Madan R, Kilambi R, Chander S. Radiation induced liver disease: a clinical update. *J Egypt Natl Canc Inst*. 2016;28(1):7–11.
46. Nazaroff WW. Radon transport from soil to air. *Rev Geophys*. 1992;30(2):137–160.
47. Hassan NM, Hosoda M, Ishikawa T, Sorimachi A, Sahoo SK, Tokonami S, et al. Radon migration process and its influence factors; review. *Jpn J Health Phys*. 2009;44(2):218–231.
48. United Nations Scientific Committee on the Effects of Atomic Radiation. Sources and effects of ionizing radiation (Volume 1). New York, NY: United Nations Scientific Committee on the Effects of Atomic Radiation; 2000.