Comparison of Skin Injury Induced by β - and γ -irradiation in the Minipig Model

Joong-Sun Kim^{1,2}, Hyosun Jang¹, Min-Ji Bae², Sehwan Shim¹, Won-Seok Jang¹, Sun-Joo Lee¹, Sunhoo Park¹, Seung-Sook Lee^{1,*}

¹Lab. of Radiation Exposure and Therapeutics, Korea Institute of Radiological and Medical Sciences, Seoul, Korea; ²Research Center, Dongnam Institute of Radiological & Medical Sciences, Busan, Korea

ABSTRACT

Background: The effects of radiation on tissues vary depending on the radiation type. In this study, a minipig model was used to compare the effects of β -rays from ^{166}Ho and γ -rays from ^{60}Co on the skin.

Materials and Methods: In this study, the detrimental effects of β - and γ -irradiation on the skin were assessed in minipigs. The histopathological changes in the skin from 1 to 12 weeks after exposure to 50 Gy of either β - (using ^{166}Ho patches) or γ - (using ^{60}Co) irradiation were assessed.

Results and Discussion: The skin irradiated by β -rays was shown to exhibit more severe skin injury than that irradiated by γ -rays at 1-3 weeks post-exposure; however, while the skin lesions caused by β -rays recovered after 8 weeks, the γ -irradiated skin lesions were not repaired after this time. The observed histopathological changes corresponded with gross appearance scores. Seven days post-irradiation, apoptotic cells in the basal layer were detected more frequently in β -irradiated skin than in γ -irradiated skin. The basal cell density and skin thickness gradually decreased until 4 weeks after γ - and β - irradiation. In β -irradiated skin lesions, the density and thickness increased sharply back to control levels by 6-9 weeks. However, this was not the case in γ -irradiated skin lesions. In γ -irradiated skin, cyclooxygenase-2 (COX-2) was shown to be expressed in the epidermis, endothelial cells of vessels, and fibroblasts, while β -irradiated lesions exhibited COX-2 expression that was mostly limited to the epidermis.

Conclusion: In this study, β -rays were shown to induce more severe skin injury than γ -rays; however, the β -rays-induced injury was largely repaired over time, while the γ -rays-induced injury was not repaired and instead progressed to necrosis. These findings reveal the differential effects of γ - and β -irradiation on skin and demonstrate the use of minipigs as a beneficial experimental model for studying irradiation-induced skin damage.

Keywords: Minipig, β -irradiation, γ -irradiation, Skin, COX-2

Original Research

Received August 17, 2017
Revision September 21, 2017
Accepted October 10, 2017

Corresponding author: Seung-Sook Lee

Lab. of Radiation Exposure and Therapeutics, Korea Institute of Radiological & Medical Sciences, 75 Nowon-gil, Nowon-gu, Seoul 01812, Korea
Tel: +82-2-970-1268
Fax: +82-2-970-2430
E-mail: sslee@kcch.re.kr

This is an Open-Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright © 2017 The Korean Association for Radiation Protection

Introduction

The skin, comprising the epidermis and the vascularized dermis, provides a vital protective barrier against environmental insults and is one of the most complex and reactive organs in the body. The increasing use of radioactive materials in industry, medicine, science, and the military as well as in localized areas of high radiation within nuclear facilities have significantly increased the potential of large-scale, uncontrolled exposure to radiation [1]. Radiation-induced skin injury is a common side-effect of radiation exposure in therapeutic applications and is also the most common problem in radiation accidents [2]. These radiation skin toxicities can negatively affect the quality

of a patient's life in terms of the pain associated with the injury as well as the premature termination of radiation treatment [3]. Radiation-induced skin injury begins with erythema, followed as the dose increases by dry and moist desquamation. Chronic radiation skin injury also results in delayed ulcers, fibrosis, and telangiectasis [4]. Although these symptoms of radiation-induced skin damage have been described, the biological mechanism underlying these effects has not yet been studied [2, 5].

The effects of radiation on tissues vary depending on the radiation type. Gamma rays are used in a wide variety of medical applications such as CT scans and radiation therapy because they exert enough energy to penetrate tissues. Beta decay results from the emission of an electron (or negative beta particle) from a nucleus. Although β -rays have weak transmission power, they have sufficient energy to penetrate the skin. β -rays may therefore be an external radiation risk [6]. The biological effects of β -irradiation in normal tissues and tumors are of interest for clinical radiotherapy, for radiation protection purposes, as well as for a greater understanding of radiation-induced inactivation of cells for different radiation types [7]. The effects of beta radiation on the skin, however, have not been well studied.

The pig has been widely used as an experimental animal in biomedical research for many centuries. The physiological and anatomical similarities between humans and pigs have deemed this animal an appropriate model for humans in many research areas [8]. Human and porcine skin tissues have microscopically heterogeneous structures that are similar in their morphology, cellular composition, and physiological properties [9]. Porcine skin has been used in drug delivery studies, to assess wound healing after heat or chemical burning, as well as to model light distribution in the skin [10].

In this study, a minipig model was used to compare the effects of β -rays from ^{166}Ho and γ -rays from ^{60}Co on the skin.

Materials and Methods

1. Animals

Six-month-old male Göttingen minipigs (mean weight, 19 kg; range 18–20 kg), which were obtained from the PWG genetics (Seoul, Korea), were used in this experiment. Two minipigs were fed a standard animal diet. All animal experiments were carried out according to protocols approved by the Institutional Animal Care and Use Committee of the Korea Institute Radiological and Medical Sciences (KIRAMS).

2. Irradiation and sample biopsy

To assess the effects of γ - and β -rays on the minipig skin, irradiation was carried out on the flank skin of minipigs. For all procedures the animals were anesthetized with tiletamine/zolazepam (Zoletil 50®, Virak Korea, Seoul, Korea) and medetomidine (Domitol®, Pfizer Animal Health Korea). To observe the skin effect of radiation, we used the right and left front back lesion of dorsal skin. We evaluated clinical status scoring in the right flank and skin biopsy performed in the left flank. Three to four days prior to irradiation, the hair in the treatment area on each pig was clipped and the position of the fields of interest was marked and tattooed with India ink. The fields were gamma irradiated with a range of 50 Gy using a ^{60}Co γ -rays irradiation therapy unit (Theratron 780, AECL, Ontario, Canada) at a dose-rate of $130.1 \text{ cGy} \cdot \text{min}^{-1}$ (field size: 5 cm, source-to-surface distance: 80 cm, depth: 1 cm [bolus 1 cm]). Each pig received 50 Gy irradiation.

To assess the effects of β emission (^{166}Ho) on the minipig skin, irradiation was carried out on the dorsal skin of minipigs. $^{166}\text{Ho}(\text{NO}_3)_3 \cdot 5\text{H}_2\text{O}$ (231 g) and 346 mL tetrahydrofuran (Merck, München, Germany) were completely dissolved in a solvent mixture of 235 mL dimethylformamide (Merck), 2 L tetrahydrofuran (Merck), and 462 g polyurethane (Dow chemicals, Midland, MI) at room temperature. The solution was uniformly affixed to adhesive tape (approximately 500 μm in thickness). Patches of the treated tape were assailed in a nuclear reactor at a neutron flux of $1 \times 10^{13} \text{ n} \cdot \text{cm}^{-2} \cdot \text{sec}^{-1}$ to convert ^{165}Ho to ^{166}Ho , which is a β emitter ($E_{\text{max}} = 1.84 \text{ MeV}$) with a half-life of 26.9 hours. ^{166}Ho also emits gamma photons (5.4% of 0.081 MeV and 0.9% of 1.38 MeV). To assess the effects of β emission (^{166}Ho) on the minipig skin, irradiation was carried out by applying the Holmium patches on the skin of minipigs. Six pieces of $1 \times 7 \text{ cm}$ -sized rectangular patches were separately applied on the dorsal skin surface and were firmly affixed with adhesive tape for 23 minutes for a total surface radiation dose of 50 Gy. At the time of irradiation, three skin biopsies were taken and pinned to cork to maintain the original size. A 5 mm punch biopsy was taken under anesthesia from the irradiated skin area and the corresponding skin area of the healthy skin. Samples of the biopsy were processed for embedding in paraffin wax after fixation in 10% buffered formalin.

3. Examination of clinicopathological alterations in gross appearance

The skin of the minipigs was carefully evaluated for 12

weeks after irradiation and the observed skin reactions were scored using a clinical status scoring system. Irradiated lesions were assessed for skin reactions and operation scars were examined. The following scoring system was used to measure radiation acute reactions in the skin: Grade 0, normal; Grade 10-30, slight swelling, edema, and erythema; Grade 40-60, dry desquamation and mild early moist desquamation changes; Grade 70-90, severe moist desquamation with some necrosis; and Grade 100, severe necrosis and loss of epidermis [5, 13].

4. Histological examination

For analysis by light microscopy, tissue samples of the skin biopsies of each field that were fixed in 10% buffered formalin, dehydrated in grade ethanol, and embedded in paraffin were assessed. The tissue sections were cut to 4 μ m thickness on a rotary microtome (Leica Ultracut, Leica Microsystems, Wetzlar, Germany), mounted on clean glass slides, cleared, hydrated, and finally stained with haematoxylin and eosin (H&E) for histological evaluation of skin injury, apoptotic cells, epidermal thickness, density of basal cells, and density of epithelial cells according to standard protocols. The slides were labeled using codes to prevent observer bias during evaluation. All tissue sections were examined by microscopy for characterization of histopathological changes. Photographs taken from skin samples were digitized using a digital camera mounted on a microscope (Leica DM IRBE, Leica Microsystems GmbH, Wetzlar, Germany). The resulting images were analyzed using image analysis software (Leica QWin, Leica Microsystems, Wetzlar, Germany).

The longest rete ridge on each slide was measured from the bottom of the basal layer to the bottom of the stratum corneum, avoiding areas where the inclusion seemed to be oblique. Means were calculated from the measurements for each slide. The same method was adopted to measure the suprapapillary thickness.

Cell density in the basal layers was determined by counting cells along a minimum of 5 mm of basement membrane. Results were expressed as the count per millimeter length of basement membrane. The degenerate cells, i.e., those cells exhibiting pycnosis and shrinkage necrosis, were excluded from these calculations.

5. Immunohistochemical study

Fixed tissues were processed in paraffin, cut into 4 μ m-thick coronal sections, and deparaffinized. The sections were

incubated with 5% normal goat serum for 60 minutes and then with rabbit anti-mouse cyclooxygenase (COX)-2 (18-7379, 1:200 dilution; Zymed) in PBS-T overnight at 4°C. The sections were then reacted with biotinylated goat anti-rabbit IgG. Immunoreactivity was then assessed using the avidin-biotin peroxidase complex (Vectastatin Elite ABC kit). The peroxidase reaction was developed using a diaminobenzidine substrate kit (DAB Substrate Kit SK-4100; Vector Laboratories). As a control, the primary antibodies were omitted for a few test sections in each experiment. The sections were counterstained with hematoxylin before being mounted and examined by microscopy.

6. Statistical analysis

The data are reported as the mean $\pm$ standard error of the mean (SEM), and data analysis was performed using one-way analysis of variance (ANOVA) followed by a Student-Newman-Keuls post hoc test for multiple comparisons. In all cases, a p -value < 0.05 was considered significant.

Results and Discussion

Time-dependent changes in skin reactions after β - and γ -irradiation of 50 Gy are summarized in Figure 1. The range of acute change begins with erythema. Over time, pigmentation, epilation, and desquamation appear in irradiated skin lesion. Between the two radiation types, β -irradiation was shown to exhibit more severe skin injury (Figure 1A and 1B) than γ -irradiation at 1-4 weeks after irradiation (Figure 1E and 1F). At eight weeks post-irradiation, the β -irradiated skin lesion had gradually repaired (Figure 1C and 1D); however, the γ -irradiated skin had not recovered and progressed to necrosis (Figure 1G and 1H).

Apoptotic cells were easily recognized based on the condensation of the cytoplasm and nuclear chromatin in these cells. Dead cells break up into several fragments and not all fragments necessarily contain nuclear material. These cytoplasmic fragments can usually be recognized by their eosinophilic staining properties in H&E staining [11]. The apoptotic cell fraction was 2-fold higher in the β -irradiated skin compared with the γ -irradiated skin at 7 days post-irradiation (Figure 2D) and apoptotic cells were mainly detected in the basal layer compared with the granular layer. A small number of cells in the basal layer exhibited apoptosis and the maximal frequency was observed at 7 days after irradiation (Figure 2A-2C). In H&E staining, demonstrated β -rays-

Fig. 1. Clinical changes in the skin of minipig following acute β - and γ -irradiation of 50 Gy. (A) 1 week, (B) 4 weeks, (C) 8 weeks, and (D) 12 weeks after β -irradiation; (E) 1 week, (F) 4 weeks, (G) 8 weeks, and (H) 12 weeks after γ -irradiation; (I) Time-dependent alteration of scoring of skin reaction in the skin of pig following β - and γ -irradiation.

Fig. 2. Alteration of apoptotic cell in the skin of minipig following acute β - and γ -irradiation of 50 Gy. (A) Normal control, (B) β -irradiation, and (C) γ -irradiation of the skin at 7 days after radiation exposure. Apoptotic cells (arrow heads) were detected in the basal cell layer of the skin by hematoxylin and eosin staining; (D) The number of apoptotic cells in the skin of minipigs at 7 days following β - and γ -irradiation.

induced desquamation of the epidermis until 4 weeks post-irradiation; however, this effect was not observed in the dermal layer (Figure 3A-3D). Epidermal detachment and accumulation of inflammatory cells in the dermis of gamma-irradiated lesions were observed (Figure 3E-3H) and the alterations in basal cell density and epidermal depth were in accordance with gross changes (Figure 3I and 3J). β -irradiation-

induced skin lesions exhibited diminished basal cell density and epidermal depth earlier than the lesions induced by γ -irradiation at 1-4 weeks after radiation; however, basal cell density and epidermal depth in the β -irradiated skin increased sharply to higher than control levels at 6-8 weeks post-irradiation and then gradually returned to control levels thereafter. The γ -irradiated skin, on the other hand, did not

Fig. 3. Histological changes in the skin of minipig following β - and γ -irradiation of 50 Gy. (A) 1 week, (B) 4 weeks, (C) 8 weeks, and (D) 12 weeks after β -irradiation; (E) 1 week, (F) 4 weeks, (G) 8 weeks, and (H) 12 weeks after γ -irradiation; Hematoxylin and eosin (400 $\times$); (I) Time-dependent changes of basal cell density in the skin of minipig following β - and γ -irradiation; (J) Time-dependent changes of epidermis depth in the skin of minipig following β - and γ -irradiation.

exhibit signs of repair (Figure 3C, 3D, 3G, and 3H).

In normal pig skin, COX-2 staining was shown to be minimal with some staining in the sebaceous glands and subcutaneous tissue, while there was no staining in the epidermis (Figure 4A). In the γ -irradiated skin, COX-2 was expressed not only in the epidermis but also in endothelial cells and fibroblasts. This effect increased with increasing severity of damage (Figure 4E-4H). The γ -irradiated skin lesions were characterized by the expression of COX-2 in the vessels and nerves of the dermis and this staining pattern increased in over time. In the β -irradiated samples, transient COX-2 expression was only observed in the epidermis (Figure 4A-4D). By 1 week post-irradiation, the expression of COX-2 was mostly limited to the stratum granulosum in the epidermis (Figure 4A). By weeks 6-10, increased expression of COX-2 was observed in all epidermal layers (Figure 4C) and was particularly prominent in the basal layer underlying areas of epidermal hyperplasia. The expression of COX-2 was shown

to decrease as skin injury recovered at 10-12 weeks after irradiation (Figure 4D and Table 1).

^{166}Ho may be an ideal radioisotope as it emits high-energy β -rays ($E_{\text{max}} = 1.84 \text{ MeV}$) with a maximum soft tissue range of 8.7 mm. As much as 90% of the β -energy from ^{166}Ho is deposited within 2.1 mm of the skin (upper and mid dermal level of the skin) [12]. This rapid dose reduction over penetration distance would result in significantly higher doses being applied to the cells in the upper viable layer of the epidermis than to the basal layer. In particular the potential stem cells, which are largely distributed towards the bases of the rete ridges, would receive far lower doses of radiation than the cells in the upper layers [3].

The time-course of the development of gross changes in response to irradiation as well as the differences in these time courses between the different irradiation types reflect the evolution and time course of changes in the component populations of the epidermis, dermis, and microvasculature

Fig. 4. COX-2 expression in the skin of minipig following β - and γ -irradiation of 50 Gy. COX-2 expression of the skin at (A) 1 week, (B) 4 weeks, (C) 8 weeks, and (D) 12 weeks after β -irradiation; COX-2 expression of the skin at (E) 1 week, (F) 4 weeks, (G) 8 weeks, and (H) 12 weeks after γ -irradiation (400 $\times$).

Table 1. Immunohistochemical Staining for COX-2 in Various Epidermal Layers of Pig Skin after Treatment with 50 Gy of Gamma or Beta ray

	Radiation	Sham	7 d	14 d	21 d	28 d	35 d
Beta-ray	Epidermis	-	+	+	++	+++	++
	Basal cell	-	+	+	++	+++	++
	Fibroblast	-	-	-	-	-	-
	Vessel	-	-	-	-	-	-
	Lipids	-	-	-	-	-	-
	Hair follicle	+	+	+	+	+	+
	Gland	-	-	-	-	-	-
Gamma-ray	Epidermis	-	+	++	+++	+++	
	Basal cell	-	++	+++	+++	+++	
	Fibroblast	-	+	++	++	++	+++
	Vessel	-	++	+++	+++	+++	+++
	Lipids	-	++	++	++	++	++
	Hair follicle	+	+	++	++	++	++
	Gland	-	+	++	+++	+++	+++

Immunolabeled sections were scored for the density of positive cells per field as follows: -, no immunoreactivity; +, weak; ++, moderate; +++, intense immunoreactivity.

of the skin [5]. The range of acute changes begins with erythema and as time passes, pigmentation, epilation, and desquamation appear in the skin lesions. The skin was usually found to develop erythema within 1 week after β -rays treatment and this reaction worsened until 4 weeks post-irradiation as evidenced by dusky/mauve erythema. After 4 weeks post- β -rays irradiation, the clinical symptoms gradually disappeared and recovered [13]. The γ -ray effects were found to be similar to the β -ray effects in the early stages after exposure; however, γ -irradiation-induced injury was not repaired and instead progressed to necrosis [13-15].

The application of β - or γ -rays to skin generally appears to trigger a linear loss of basal cells [13-16] and histological studies have demonstrated an early pyknosis of cells in the

basal layers of the epidermis 7 days after irradiation, which in turn initiates an apoptotic reaction and the subsequent disruption of the total epidermis. In a previous study we showed that apoptotic cells were present at a higher frequency in the basal layer of irradiated skin than in the basal layer of normal skin and that the frequency was irradiation dose-dependent [15]. In this study, we showed that β -irradiated skin contains a higher proportion of pyknotic cells than γ -irradiated skin.

Changes in basal cell density and epidermal depth were previously shown to correspond to gross skin morphology scores [13-16]. Epidermal thickness began to decrease 1 week after radiation-induced skin injury, which was evident 6-8 weeks after β - or γ -irradiation. The β -rays have a faster onset of erythema and moist desquamation after exposure

to radiation, while beta radiation- and not gamma radiation-induced injury was followed by a pronounced degree of recovery. Gamma-rays penetrate the whole skin layer, causing damage to blood vessels and lipids in the dermis and epidermal layer. This damage leads to erosion and ulcers, where the disappearance of the epidermal layer results from the invasion of secondary acute inflammatory cells [15].

The differences in recovery patterns between the two radiation types relates to interactions between keratinocytes and fibroblasts: keratinocytes stimulate fibroblasts to synthesize growth factors such as keratinocyte growth factor, hepatocyte growth factor, and transforming growth factor beta, which in turn stimulate keratinocyte proliferation in a double paracrine manner [17]. Accordingly, γ -irradiation-induced injury was delayed due to the damage to the dermal layer containing fibroblasts and endothelial cells disrupting in the interaction of keratinocytes and fibroblasts.

In this study, sequential skin tissue samples were obtained from the same individual, which allowed for continuous clinicopathological changes including the potential role of COX-2 in radiation-induced skin damage to be monitored. COX-2, a stress response protein, is known to produce the synthesis of prostaglandins involved in the inflammatory cascade resulting in tissue injury [18, 19]. In radiation-associated mucositis, the role of COX-2 has been reported to increase the severity of mucosal injury and to prolong the inflammatory and ulcerative event [19]. COX-2 expression in the epidermis is furthermore known to be induced by stimuli that result in transient hyperplasia, such as UV irradiation, including in UV-induced skin cancers and other malignancies [17]. The expression of COX-2 in radiation-induced skin damage due to high-dose ionizing irradiation has not yet been reported. Although this study does not confirm the role of COX-2 in the irradiated skin, COX-2 expression was shown here to be one of the irradiation-induced skin injury markers. In this study, COX-2 was shown to be expressed not only in the epidermis but also in endothelial cells and in fibroblasts in the γ -irradiated skin, where the expression was shown to increase with increasing severity of damage. In this study, γ -irradiation-induced skin lesions were characterized by COX-2 expression in the vessels and nerves of the dermis and lipids and the γ -irradiated skin of minipigs were shown to exhibit increasing COX-2 expression over time. The skin of minipigs irradiated with β -rays was shown to exhibit only transient COX-2 expression and only in the epidermis.

Conclusion

In this study, the biological effects of β - and γ -irradiation on the skin of minipigs were assessed and compared. Overall, β -rays were shown to induce more severe skin injury than γ -rays; however, the β -rays-induced injury was largely repaired over time, while the γ -rays-induced injury was not repaired and instead progressed to necrosis. These findings suggest that γ -irradiation affects the deep dermis and subcutaneous fat due to its penetrating ability, whereas β -rays irradiation induces changes only at the level of the epidermis and the upper dermis. The use of the minipig in this study demonstrates the use of the animal model as an experimental human skin model with which radiation-induced skin damage can be studied and thus as a model with which the efficacy of therapeutics to reduce the skin damage and to accelerate repair can be developed.

Acknowledgements

This study was supported by the Nuclear R&D Program of the Ministry of Education, Science and Technology, Korea (50581-2016).

References

1. Wang XJ, et al. The effect of RHIZOMA COPTIDIS and COPTIS CHINENSIS aqueous extract on radiation-induced skin injury in a rat model. *BMC Complementary and Altern. Med.* 2013;15:13:105.
2. Hopewell JW. The skin: its structure and response to ionizing radiation. *Int. J. Radiat. Biol.* 1990;57:751-773.
3. McQuestion M. Evidence-based skin care management in radiation therapy: clinical update. *Semin. Oncol. Nurs.* 2011;27:e1-17.
4. Mendelsohn FA, Divino CM, Reis ED, Kerstein MD. Wound care after radiation therapy. *Adv. Skin and Wound Care.* 2002;15:216-224.
5. Archambeau JO, Ines A, Fajardo LF. Pathophysiology of irradiated skin and breast. *Int. J. Radiat. Oncol. Biol. Phys.* 1995;31:1171-1185.
6. Peel DM, Hopewell JW, Wells J, Charles MW. Nonstochastic effects of different energy beta emitters on pig skin. *Radiat. Res.* 1984;99:372-382.
7. Chung YL, Lee JD, Bang D, Lee JB, Park KB, Lee MG. Treatment of Bowen's disease with a specially designed radioactive skin patch. *Eur. J. Nucl. Med.* 2000;27:842-846.
8. Swindle MM, Makin A, Herron AJ, Clubb FJ Jr, Frazier KS. Swine

- as models in biomedical research and toxicology testing. *Vet. Pathol.* 2012;49:344-356.
9. Sullivan TP, Eaglstein WH, Davis SC, Mertz P. The pig as a model for human wound healing. *Wound Repair Regener.* 2001;9:66-76.
10. Ma ZF, Chai JK, Yang HM, Liu Q, Xu MH, Yin HN. Acellular porcine dermal matrix produced with different methods and an experimental study on its transplantation to skin wound. *Chinese critical care medicine.* 2005;17:92-94.
11. Martin P. Wound healing-aiming for perfect skin regeneration. *Science.* 1997;276:75-81.
12. Johnson LS, Yanch JC, Shortkroff S, Barnes CL, Spitzer AI, Sledge CB. Beta-particle dosimetry in radiation synovectomy. *Eur. J. Nucl. Med.* 1995;22:977-988.
13. Kim JS, Rhim KJ, Jang WS, Lee SJ, Son Y, Lee SS, Park S, Lim SM. β -irradiation (^{166}Ho patch)-induced skin injury in mini-pigs: effects on NF- κ B and COX-2 expression in the skin. *J. Vet. Sci.* 2015;16:1-9.
14. Kim JS, Jang WS, Bae MJ, Park S, Lee SS. Radiation-induced eosinophilic, polymorphic, and pruritic eruption in a pig skin model. *Lab. Anim. Res.* 2015;31:204-208.
15. Kim JS, Park SH, Jang WS, Lee SJ, Lee SS. Gamma-ray-induced skin injury in the mini-pig: Effects of irradiation exposure on cyclooxygenase-2 expression in the skin. *J. Radiat. Protect. Res.* 2015;40:65-72.
16. Kim J, Park S, Jeon BS, Jang WS, Lee SJ, Son Y, Rhim KJ, Lee SI, Lee SS. Therapeutic effect of topical application of curcumin during treatment of radiation burns in a mini-pig model. *J. Vet. Sci.* 2016;17:435-444.
17. Werner S, Krieg T, Smola H. Keratinocyte-fibroblast interactions in wound healing. *J. Invest. Dermatol.* 2007;127:998-1008.
18. Tripp CS, Blomme EA, Chinn KS, Hardy MM, LaCelle P, Pentland AP. Epidermal COX-2 induction following ultraviolet irradiation: suggested mechanism for the role of COX-2 inhibition in photoprotection. *J. Invest. Dermatol.* 2003;121:853-861.
19. Yeoh AS, Bowen JM, Gibson RJ, Keefe DM. Nuclear factor kappaB (NFkappaB) and cyclooxygenase-2 (Cox-2) expression in the irradiated colorectum is associated with subsequent histopathological changes. *Int. J. Radiat. Oncol. Biol. Phys.* 2005;63:1295-1303.